

PROPHYLACTIC PLAN FOR THE GUATEMALAN ARMY

I. Educational Program

(a) Teach departmental physicians and physicians in charge of each barrack.

(b) Physicians teach all of the soldiers

(c) Teaching methods

1. Movies
2. Talks
3. Posters

2. Current prophylaxis

(a) Materials

1. Condom
2. Silver proteinate solution -fresh  
10% argyrol  
2% protargol
3. Ointment

	Parts
Hydrargyri chloridum mite	30
Adeps benzoinatus	65
Cere alba U.S.P	5

(b) Methods


1. The soldier receives a condom before leaving the barracks.
2. Upon his return he must go to the Department of Prophylaxis where he will receive treatment. The treatment needs to be monitored by a well-trained assistant.

(c) Routine

1. Urinate as soon as he arrives at the barracks.
2. Wash well with soap and water, preferably using tincture of green soap, for 5 minutes. The penis with the foreskin pulled back, the scrotum and all of the surrounding skin need to be washed thoroughly.
3. Inject 4 cc of silver proteinate solution. The injected solution must be retained for 5 minutes allowing drops to occasionally come out.
4. Rub in well 2 to 4 grams of Calomel ointment on the penis, scrotum and surrounding skin.
5. Cover all of the treated area well with paper to avoid staining clothes. Patient must wait 5 hours before urinating.

[TN: Page 2 is a copy of page 1. Everything is the same.]

I, Jiri Stejskal, Ph.D., hereby certify that the above document is an accurate translation from Spanish into English.  
Date: July 20, 2011

Signature: 

CETRA, Inc.  
7804 Montgomery Ave., Suite 8, Elkins Park, PA 19027

Tel.: (215) 635-7090

CTLR\_0001073\_ENG